

Purification and characterization of serine proteinase 2 from *Bacillus intermedius* 3-19

Balaban N., Mardanov A., Sharipova M., Gabdrakhmanova L., Sokolova E., Rudenskaya G., Leshchinskaya I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

A proteinase secreted in the late stationary phase was isolated from the culture fluid of *Bacillus intermedius* 3-19 by ion-exchange chromatography on CM-cellulose followed by FPLC on a Mono S column. The enzyme was completely inhibited by the serine proteinase inhibitors diisopropyl fluorophosphate and phenylmethylsulfonyl fluoride. The maximum proteolytic activity against the synthetic chromogenic substrate Z-Ala-Ala-Leu-pNA was observed at pH 9.0. The molecular weight of the enzyme is 28 kD and its isoelectric point is 9.2. We have also determined pH- and thermostability and K_m and k_{cat} of this proteinase. The enzyme has been classified as a thiol-dependent serine proteinase. N-Terminal amino acid sequence (10 residues) and amino acid composition of the protein were also determined. By the mode of hydrolysis of peptide bonds in the oxidized B-chain of insulin, this enzyme is similar to the thiol-dependent serine proteinase 1 from *B. intermedius* 3-19 secreted during vegetative growth.

<http://dx.doi.org/10.1023/B:BIRY.0000026198.81752.f4>

Keywords

Properties, Proteinase, Purification, Thiol-dependent serine proteinase (*Bacillus intermedius*)